

ON EQUATIONS DEFINING COINCIDENT ROOT LOCI

JAYDEEP V. CHIPALKATTI

Abstract. We revisit an old problem in classical invariant theory, viz. that of giving algebraic conditions for a binary form to have linear factors with assigned multiplicities. We construct a complex of SL_2 -representations such that the desired algebraic conditions are expressible as a specific cohomology group of this complex.

Mathematics Subject Classification: 14L30, 16W22

1. INTRODUCTION

Consider a binary form $F(x, y) = \sum_{j=0}^d a_j x^j y^{d-j}$ over the complex numbers. It splits as a product of linear forms, and it is classical that F has a repeated factor if and only if its discriminant vanishes. Similarly, we can ask for algebraic conditions on the coefficients a_j , so that F has say, a triple factor or two double factors. More generally, we may fix a partition λ of d , and ask for algebraic conditions so that the factors of F have multiplicities as dictated by the parts in λ . The object of this note is to devise a method for answering such questions.

This problem is addressed for the first time (to my knowledge) by Arthur Cayley [4]. Thereafter J. Weyman has obtained substantial results for special kinds of partitions in [18], [19] and [20] (see section 6).

The polynomial $F(x, y)$ as above will be identified with the point $[a_0, \dots, a_d]$ of $\mathbb{P}^d$. Let then $\lambda = (\lambda_1, \dots, \lambda_n)$ be a partition of d , and consider the set

$$X_\lambda = \{[a_0, \dots, a_d] : F \text{ factors as } \prod_{i=1}^n L_i^{\lambda_i} \text{ for some linear forms } L_i\},$$

which is a projective subvariety of $\mathbb{P}^d$. This is the coincident root locus in the title and the principal object of study in this paper.

The group $SL_2(\mathbb{C})$ acts on the imbedding $X_\lambda \subseteq \mathbb{P}^d$ in the following manner. An element $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in SL_2(\mathbb{C})$ sends the form F to $F' = \sum_{j=0}^d a'_j x^j y^{d-j}$, where the a'_j are determined by

$$\sum_{j=0}^d a_j (\alpha x + \beta y)^j (\gamma x + \delta y)^{d-j} = \sum_{j=0}^d a'_j x^j y^{d-j}.$$

Its defining ideal $I_{X_\lambda} < \mathbb{C}[a_0, \dots, a_d]$ is then an SL_2 -subrepresentation, and our problem is essentially one of calculating this ideal. Our main result (Corollary 3.5) is a (somewhat indirect) solution to this problem. Specifically, for each positive integer m , we construct a complex of $SL_2(\mathbb{C})$ -representations (see formula (16)) whose zeroth cohomology group is the graded piece $(I_X)_m$. In section 4, we use this complex to calculate the ideal for a few specific examples and express the answer in the language of classical invariant theory (especially see Theorems 4.3 and 4.4). In Theorem 5.4 of section 5, we identify the singular locus of X_λ ; a result which should be of interest in itself.

The base field will be $\mathbb{C}$. All terminology from algebraic geometry follows Hartshorne [10]. The formulae are numbered in a single sequence throughout the text.

ACKNOWLEDGEMENTS: The program Macaulay-2 has been of immense help in computations, and it is a pleasure to thank its authors Dan Grayson and Mike Stillman. The financial assistance of Anthony V. Geramita, Leslie Roberts and Queen's University during the progress of this work is gratefully acknowledged.

2. PRELIMINARIES

In the next three subsections, we recall a few matters from the invariant theory of binary forms. An excellent reference for the classical theory is the text by Grace and Young [7]. See [6] and [16] for the modern theory and [17] for a discussion of algorithms for the computation of invariants and covariants.

2.1. Representations of $SL_2(\mathbb{C})$. In the sequel, V denotes a two dimensional $\mathbb{C}$ -vector space. We will use the following isomorphisms

of $SL(V)$ –modules :

$$(1) \quad \begin{aligned} \wedge^m(\mathrm{Sym}^n V) &= \mathrm{Sym}^m(\mathrm{Sym}^{n+1-m} V), \\ \mathrm{Sym}^m V \otimes \mathrm{Sym}^n V &= \bigoplus_{r=0}^{\lfloor \frac{m+n}{2} \rfloor} \mathrm{Sym}^{m+n-2r} V \quad (\text{Clebsch–Gordan formula}); \end{aligned}$$

and the Cayley–Sylvester formula

$$(2) \quad \mathrm{Sym}^m(\mathrm{Sym}^n V) = \bigoplus_{r=0}^{\lfloor \frac{mn}{2} \rfloor} (\mathrm{Sym}^{mn-2r} V)^{\oplus p(r,m,n)-p(r-1,m,n)},$$

where $p(r, m, n)$ is the number of partitions of r into at most m parts with no part exceeding n . The $GL(V)$ –modules $\mathrm{Sym}^d(V^*)$ and $(\mathrm{Sym}^d V)^*$ are isomorphic, and we write them indifferently as $\mathrm{Sym}^d V^*$.

Let $\{x, y\}$ be a basis of V . With the identification $a_j = (x^j y^{d-j})^* \in \mathrm{Sym}^d V^*$, the form $F = \sum a_j x^j y^{d-j}$ corresponds to the trace element $\sum_j (x^j y^{d-j})^* \otimes x^j y^{d-j}$ in $\mathrm{Sym}^d V^* \otimes \mathrm{Sym}^d V$. The ring $\mathbb{C}[a_0, \dots, a_d]$ is identified with the symmetric algebra $\mathrm{Sym}^\bullet(\mathrm{Sym}^d V^*)$.

2.2. Invariants and Covariants. Let

$$\mathrm{Sym}^q V^* \hookrightarrow \mathrm{Sym}^p(\mathrm{Sym}^d V^*)$$

be an $SL(V)$ –subrepresentation; or what is the same, a nonzero $SL(V)$ –morphism $\mathbb{C} \longrightarrow \mathrm{Sym}^p(\mathrm{Sym}^d V^*) \otimes \mathrm{Sym}^q V$. Its image consists of all scalar multiples of a polynomial $\Phi(a_0, \dots, a_d; x, y)$, having total degree p in the a_j and q in x, y . This polynomial is called a covariant of degree p and order q (of the binary form F). We will also say that Φ is of type (p, q) . E.g., for $d \geq 2$, the inclusion $\mathrm{Sym}^{2d-4} V^* \hookrightarrow \mathrm{Sym}^2(\mathrm{Sym}^d V^*)$ defines a covariant of type $(2, 2d-4)$, called the Hessian of F .

An invariant is a covariant of order zero. E.g., the discriminant of F (defined to be the Sylvester resultant of $\partial F/\partial x$ and $\partial F/\partial y$) is an invariant of degree $2d-2$.

2.3. Transvectants. Let Φ_1, Φ_2 be covariants of types $(p_1, q_1), (p_2, q_2)$ respectively, and $1 \leq r \leq \min\{q_1, q_2\}$ an integer. Associated to these is a covariant of type $(p_1 + p_2, q_1 + q_2 - 2r)$ called the r -th transvectant of Φ_1, Φ_2 , written $(\Phi_1, \Phi_2)^r$. (By convention (Φ_1, Φ_2) stands for $(\Phi_1, \Phi_2)^1$.) For instance, the Hessian equals $(F, F)^2$.

Scholium 2.1. The transvectant can be defined in several ways, and the various definitions are in agreement only upto a scalar. In section 4, it will be necessary for us to fix this scalar, and it is important that some definite convention be followed consistently.

We will use the definition given in terms of the symbolic method.

Let¹ $F = \sum_{j=0}^d \binom{d}{j} a_j x^j y^{d-j}$, and

$$\Phi_i = \sum_{k=0}^{q_i} \varphi_{ik} x^k y^{q_i-k}, \quad i = 1, 2.$$

Thus each φ_{ik} is a homogeneous form of degree p_i in the a_j .

Now formally write $\Phi_1 = (\alpha_1 x + \alpha_2 y)^{q_1}$, $\Phi_2 = (\beta_1 x + \beta_2 y)^{q_2}$ and define

$$(\Phi_1, \Phi_2)^r = (\alpha_1 \beta_2 - \alpha_2 \beta_1)^r (\alpha_1 x + \alpha_2 y)^{q_1-r} (\beta_1 x + \beta_2 y)^{q_2-r}.$$

That is to say, expand the right hand side and substitute φ_{1k} and φ_{2k} respectively, for $\binom{q_1}{k} \alpha_1^k \alpha_2^{q_1-k}$ and $\binom{q_2}{k} \beta_1^k \beta_2^{q_2-k}$. The symbolic method is more sound than it appears prima facie, see [12] for a thorough modern treatment.

2.4. Definition of the Coincident Root Locus. In the sequel $\lambda = (1^{e_1} 2^{e_2} \dots d^{e_d})$ denotes a partition of d , having e_r parts of size r for $1 \leq r \leq d$. The number of parts is $\sum e_r = n$.

With $\{x, y\}$ a basis of V , we identify a point of $\mathbb{P} \text{Sym}^{e_r} V$ with a degree e_r polynomial $G_r(x, y)$ determined upto scalars.

Now we have Veronese maps

$$(3) \quad v_r : \mathbb{P} \text{Sym}^{e_r} V \longrightarrow \mathbb{P} \text{Sym}^{r e_r} V, \quad G_r \longrightarrow G_r^r$$

and a multiplication map

$$(4) \quad \mu : \prod_{r=1}^d \mathbb{P} \text{Sym}^{r e_r} V \longrightarrow \mathbb{P} \text{Sym}^d V, \quad (H_1, \dots, H_d) \longrightarrow \prod_{r=1}^d H_r.$$

¹The introduction of binomial coefficients is a familiar device in order to make things work smoother.

Finally consider the composite f_λ ,

$$(5) \quad \begin{aligned} (Y_\lambda =) \prod_r \mathbb{P} \operatorname{Sym}^{e_r} V &\xrightarrow{\Pi v_r} \prod_r \mathbb{P} \operatorname{Sym}^{r e_r} V \xrightarrow{\mu} \mathbb{P} \operatorname{Sym}^d V, \\ (G_1, \dots, G_d) &\xrightarrow{f_\lambda} \prod_{r=1}^d G_r^r. \end{aligned}$$

Definition 2.2. The Coincident Root locus X_λ is defined to be the scheme-theoretic image of f_λ .

The locus is closed in $\mathbb{P} \operatorname{Sym}^d V$ since f_λ is a projective morphism, and it is reduced since Y_λ is. Its dimension is the number of parts in λ , which is n .

The morphism $Y_\lambda \xrightarrow{f_\lambda} X_\lambda$ is finite (being quasi-finite and projective) and birational (since for a general F in the image, G_r can be recovered as the product of linear forms dividing F exactly r times). The ideal sheaf $\mathcal{I}_{X_\lambda}$ is the kernel

$$(6) \quad 0 \longrightarrow \mathcal{I}_{X_\lambda} \longrightarrow \mathcal{O}_{\mathbb{P}^d} \longrightarrow \mathcal{O}_{X_\lambda} \longrightarrow 0,$$

and for later use we define $\mathcal{D}_\lambda$ to be the cokernel

$$(7) \quad 0 \longrightarrow \mathcal{O}_{X_\lambda} \longrightarrow f_{\lambda*} \mathcal{O}_{Y_\lambda} \longrightarrow \mathcal{D}_\lambda \longrightarrow 0.$$

We will drop the suffix λ when it is safe to do so.

Scholium 2.3. • We have $X_{\lambda_1} \subseteq X_{\lambda_2}$ exactly when λ_2 is a refinement of λ_1 .

- The locus $X_{(d)}$ is the rational normal curve of degree d and $X_{(d-1,1)}$ its tangential developable. In general $X_{(d-s,1^s)}$ is the tangential developable (i.e., the closure of the union of tangent spaces at smooth points) of $X_{(d-s+1,1^{s-1})}$. Of course, $X_{(2,1^{d-2})}$ is the hypersurface of degree $2(d-1)$ defined by the discriminant.
- The degree of the CR locus in $\mathbb{P}^d$ is given by

$$\deg X_\lambda = \frac{n!}{\prod_r (e_r!)} \prod_r r^{e_r},$$

a formula going back to Hilbert. To see this, identify $\mathbb{P} \operatorname{Sym}^d V \cong \operatorname{Sym}^d(\mathbb{P}^1)$ with the set of effective divisors of degree d on $\mathbb{P}^1$. Let Σ denote a general $(d-n)$ -dimensional linear subspace of $\mathbb{P}^d$, it corresponds to a linear series g_{d-n}^d on $\mathbb{P}^1$. The points of $\Sigma \cap X$

correspond to those divisors in the series which may be written as $\sum_r r(P_{r,1} + P_{r,2} + \cdots + P_{r,e_r})$, for some points $P_{r,j} \in \mathbb{P}^1$. According to De Jonquières' formula (see [2, p. 359]), the number of such divisors is the coefficient of $t_1^{e_1} t_2^{e_2} \cdots t_d^{e_d}$ in $(1+t_1+2t_2+\cdots+dt_d)^n$, hence the assertion.

3. THE IDEAL OF THE CR LOCUS

3.1. The Machine Computation of I_X . There is a rather straightforward algorithm for the computation of the ideal I_X , which can be implemented on any computer algebra system with modest capabilities. Since X is given as the image of a projective morphism, the problem is one of elimination theory. We illustrate with the case $\lambda = (3\ 2\ 2)$. Consider the morphism

$$f_{322} : \mathbb{P} \operatorname{Sym}^2 V \times \mathbb{P} V \longrightarrow \mathbb{P} \operatorname{Sym}^7 V, \quad (G_2, G_3) \longrightarrow G_2^2 G_3^3 (= F).$$

Using affine cöordinates, write

$$G_2 = t^2 + u_1 t + u_2, \quad G_3 = t + v_1, \quad F = t^7 + a_1 t^6 + \cdots + a_7.$$

By forcing the equality

$$t^7 + a_1 t^6 + a_2 t^5 + \cdots + a_7 = (t^2 + u_1 t + u_2)^2 (t + v_1)^3,$$

we get polynomial expressions $a_j = q_j(u_1, u_2, v_1)$ for $1 \leq j \leq 7$. This defines a homomorphism

$$q_{322} : \mathbb{C}[a_1, \dots, a_7] \longrightarrow \mathbb{C}[u_1, u_2, v_1],$$

which is the ring theoretic counterpart of f_{322} . Calculate a Gröbner basis for the kernel of q_{322} (with respect to a degree lex or degree reverse-lex term order) and homogenize with respect to a new variable a_0 ; this gives the ideal I_X . (See e.g. [1, Exer. 1.6.19c].)

The actual calculation shows that it is generated by a 364-dimensional subspace of $\mathbb{C}[a_0, \dots, a_7]_6$. This is not very informative by itself; and to gain some insight into the structure of I_X , we aim to describe it invariant-theoretically. In the present example, this means identifying the subrepresentation $(I_X)_6 \subseteq \operatorname{Sym}^6(\operatorname{Sym}^7 V^*)$. To this end, we construct our main technical artifice (the complex $\mathcal{G}^\bullet$) in the next subsection.

The idea, in a nutshell, is to approach X indirectly by working with its normalisation Y . We will construct a locally free resolution of the

structure sheaf $\mathcal{O}_Y$, and then generate a spectral sequence by taking its pushforward.

3.2. The Eagon–Northcott Complex. Set $T = Y \times \mathbb{P} \operatorname{Sym}^d V$ with projections $\pi_r : T \rightarrow \mathbb{P} \operatorname{Sym}^{e_r} V$ for $1 \leq r \leq d$, and $\pi : T \rightarrow \mathbb{P} \operatorname{Sym}^d V$. Factor the map f as

$$Y \xrightarrow{1 \times f} T \xrightarrow{\pi} \mathbb{P} \operatorname{Sym}^d V,$$

and let $\Gamma \subseteq T$ denote the image of $1 \times f$.

$$\begin{array}{ccc} T & \xrightarrow{\pi_r} & \mathbb{P} \operatorname{Sym}^{e_r} V \\ \downarrow \pi & & \\ \mathbb{P} \operatorname{Sym}^d V & & \end{array} \quad \begin{array}{ccccc} Y & \xrightarrow{\text{iso}} & \Gamma & \xrightarrow{\text{incl}} & T \\ & \searrow f & \downarrow \pi & & \downarrow \pi \\ & & X & \xrightarrow{\text{incl}} & \mathbb{P}^d \end{array}$$

We claim that the structure sheaf $\mathcal{O}_\Gamma$ is resolved by an Eagon–Northcott complex of vector bundles on T .

Quite generally, let $\mathcal{A} \xrightarrow{\varphi} \mathcal{B}$ be a morphism of vector bundles on a regular scheme T . Assume that the bundles have ranks a, b respectively, with $a \geq b$. Then we have a complex

$$0 \rightarrow \mathcal{N}^{b-a-1} \rightarrow \dots \rightarrow \mathcal{N}^p \rightarrow \mathcal{N}^{p+1} \rightarrow \dots \rightarrow \mathcal{N}^0 (= \mathcal{O}_T) \rightarrow 0,$$

where

(8)

$$\mathcal{N}^p = (\wedge^b \mathcal{B})^{-1} \otimes \operatorname{Sym}^{-(p+1)} \mathcal{B}^* \otimes \wedge^{b-p-1} \mathcal{A} \quad \text{for } b-a-1 \leq p \leq -1.$$

The differential of the complex is defined via contraction with the section $\varphi \in H^0(T, \mathcal{A}^* \otimes \mathcal{B})$. (See [5, Appendix A2.6] for details of the construction.)

Let T_φ be the closed subscheme of T defined by the ideal sheaf $\operatorname{Fitt}_0(\operatorname{coker} \varphi)$. Set-theoretically, it is the *degeneracy locus* $\{t \in T : \operatorname{rank} \varphi_t \leq b-1\}$. If the subscheme has codimension $a-b+1$ and is a local complete intersection, then $\mathcal{N}^\bullet$ is a resolution of its structure sheaf $\mathcal{O}_{T_\varphi}$ (loc.cit.).

Returning to our setup, define line bundles

$$\mathcal{L} = \bigotimes_{r=1}^d \pi_r^* \mathcal{O}_{\mathbb{P}^{e_r}}(r), \quad \mathcal{M} = \pi^* \mathcal{O}_{\mathbb{P}^d}(1)$$

on T . By the Künneth formula,

$$H^0(T, \mathcal{L}) = \bigotimes_{r=1}^d \operatorname{Sym}^r(\operatorname{Sym}^{e_r} V^*), \quad H^0(T, \mathcal{M}) = \operatorname{Sym}^d V^*.$$

Define the *comultiplication* map

$$(9) \quad \mathrm{Sym}^d V^* \longrightarrow \bigotimes_{r=1}^d \mathrm{Sym}^r(\mathrm{Sym}^{e_r} V^*),$$

as dual to the usual multiplication map on polynomials. Adding it to the identity $\mathrm{Sym}^d V^* \longrightarrow H^0(T, \mathcal{M})$ gives a map

$$\mathrm{Sym}^d V^* \longrightarrow H^0(T, \mathcal{L} \oplus \mathcal{M}),$$

which defines a morphism of bundles

$$\varphi : \underbrace{\mathrm{Sym}^d V^* \otimes \mathcal{O}_T}_{\mathcal{A}} \longrightarrow \underbrace{\mathcal{L} \oplus \mathcal{M}}_{\mathcal{B}}, \quad \text{with } a = d + 1, b = 2.$$

Theorem 3.1. *The Eagon–Northcott complex of φ resolves $\mathcal{O}_\Gamma$.*

PROOF. We will show that the locus $\{\mathrm{rank} \varphi \leq 1\}$ coincides with Γ . Since Γ (being smooth) is a local complete intersection of codimension d in T , the theorem will follow.

It is easier to work with the transpose map

$$\varphi^{\mathrm{trans}} : \mathcal{L}^{-1} \oplus \mathcal{M}^{-1} \longrightarrow \mathrm{Sym}^d V \otimes \mathcal{O}_T,$$

which has the same degeneracy locus. Let then $t = (G_1, \dots, G_d; F)$ be a point of T . The fibre of the line bundle $\mathcal{O}_{\mathbb{P}^{e_r}}(-r)$ over the point $G_r \in \mathbb{P} \mathrm{Sym}^{e_r} V$ is naturally isomorphic to the linear span of G_r , hence the fibre of $\mathcal{L}^{-1}$ over t is the space $\mathbb{C} \langle \prod_r G_r \rangle$. So over t , the map φ^{trans} is given by

$$\mathbb{C} \langle \prod_r G_r \rangle \oplus \mathbb{C} \langle F \rangle \longrightarrow \mathbb{C} \langle F, \prod_r G_r \rangle.$$

This map has rank one iff F is a scalar multiple of $\prod_r G_r$ iff t lies in Γ .

The theorem is proved. $\square$

After expanding (8),

(10)

$$\mathcal{N}^p = \left\{ \bigoplus_{\substack{\alpha + \beta = -(p+1), \\ \alpha, \beta \geq 0}} \mathcal{L}^{-(\alpha+1)} \otimes \mathcal{M}^{-(\beta+1)} \right\} \otimes \wedge^{1-p}(\mathrm{Sym}^d V^*)$$

for $-d \leq p \leq -1$,

$$\mathcal{N}^0 = \mathcal{O}_T \quad \text{and} \quad \mathcal{N}^\bullet \xrightarrow{\sim}_q \mathcal{O}_\Gamma.$$

Now consider the second quadrant spectral sequence

$$(11) \quad \begin{aligned} E_1^{p,q} &= \mathbf{R}^q \pi_* \mathcal{N}^p \quad \text{for } -d \leq p \leq 0 \text{ and } q \geq 0; \\ d_r^{p,q} : E_r^{p,q} &\longrightarrow E_r^{p+r, q-r+1}, \quad E_\infty^{p,q} \Rightarrow \mathbf{R}^{p+q} \pi_* \mathcal{O}_\Gamma. \end{aligned}$$

Each $E_1^{p,q}$ term can be precisely calculated using the projection formula followed by the Künneth formula. Apart from $E_1^{0,0} = \mathcal{O}_{\mathbb{P}^d}$, all nonzero terms are in the n -th row. In fact

$$E_1^{p,n} = \left\{ \bigoplus_{\substack{\alpha+\beta=-(p+1) \\ \alpha, \beta \geq 0}} \left\{ \bigotimes_r H^{e_r}(\mathbb{P}^{e_r}, \mathcal{O}_{\mathbb{P}}(-r\alpha - r)) \right\} \otimes \mathcal{O}_{\mathbb{P}^d}(-\beta - 1) \right\} \\ \otimes \wedge^{1-p}(\text{Sym}^d V^*).$$

Now $H^{e_r}(\mathbb{P}^{e_r}, \mathcal{O}_{\mathbb{P}}(-r\alpha - r)) \neq 0$ iff $r\alpha + r \geq e_r + 1$. Using Serre duality, we get

$$(12) \quad \begin{aligned} E_1^{p,n} &= \left\{ \bigoplus_{\substack{\alpha+\beta=-(p+1) \\ \alpha \geq M-1, \beta \geq 0}} \left\{ \bigotimes_r \text{Sym}^{r\alpha+r-e_r-1}(\text{Sym}^{e_r} V) \right\} \otimes \mathcal{O}_{\mathbb{P}^d}(-\beta - 1) \right\} \\ &\quad \otimes \wedge^{1-p}(\text{Sym}^d V^*). \end{aligned}$$

Here M denotes $\lceil \max_r \{ \frac{e_r+1}{r} \} \rceil$. Now π is a finite morphism on Γ , hence $E_\infty^{p+q} = 0$ for $p+q \neq 0$ and we have an extension

$$(13) \quad 0 \longrightarrow E_\infty^{0,0} \longrightarrow \pi_* \mathcal{O}_\Gamma \longrightarrow E_\infty^{-n,n} \longrightarrow 0.$$

Proposition 3.2. *We have*

$$E_2^{p,n} = \cdots = E_{n+1}^{p,n} = \begin{cases} \mathcal{I}_X & \text{if } p = -(n+1), \\ \mathcal{D} & \text{if } p = -n, \\ 0 & \text{otherwise.} \end{cases}$$

PROOF. All the differentials from d_2 upto d_n are zero, hence $E_2^{p,n} = \cdots = E_{n+1}^{p,n}$. If $p \neq -n, -(n+1)$, then $E_2^{p,n} = E_\infty^{p,n} = 0$. Now going back to the definition of the filtration on the abutment of a spectral sequence (see e.g. [3, Ch. III]), it follows that $E_\infty^{0,0}$ is the image of the morphism $\pi_* \mathcal{O}_T \rightarrow \pi_* \mathcal{O}_\Gamma$, which is $\mathcal{O}_X$. Hence $E_2^{-n,n} = E_\infty^{-n,n}$ equals $\text{coker}(\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_\Gamma) = \mathcal{D}$. Finally $E_2^{-(n+1),n} = \ker(E_1^{0,0} \rightarrow E_\infty^{0,0}) = \mathcal{I}_X$. $\square$

The proposition exhibits $\mathcal{I}_X$ as the middle cohomology of the complex

$$E_1^{-(n+2),n} \rightarrow E_1^{-(n+1),n} \rightarrow E_1^{-n,n}.$$

Each term of the complex is a vector bundle on $\mathbb{P}^d$, explicitly described in terms of the combinatorial datum λ .

Our next step is to work with the complex given by the n -th row of this spectral sequence. Let $\mathcal{G}^\bullet$ be the complex $E_1^{\bullet, n}[-(n+1)]$, with differential denoted d_G . By the previous proposition, $\mathcal{H}^p(\mathcal{G}^\bullet) = \mathcal{I}_X, \mathcal{D}$ for $p = 0, 1$ respectively and zero elsewhere. The point of introducing the shift is merely to simplify the indices. $\mathcal{G}^p$ is nonzero in the range $n+1-d \leq p \leq n+1-M$.

3.3. Hypercohomology of $\mathcal{G}^\bullet$. Set $\mathcal{G}^\bullet(m) = \mathcal{G}^\bullet \otimes \mathcal{O}_{\mathbb{P}^d}(m)$ for $m \geq 0$. Our interest lies in the hypercohomology of the complex $\mathcal{G}^\bullet(m)$. There are two spectral sequences

$$(14) \quad E_1^{p,q} = H^q(\mathbb{P}^d, \mathcal{G}^p(m)), \quad d_r^{p,q} : E_r^{p,q} \longrightarrow E_r^{p+r, q-r+1},$$

$$(15) \quad \dot{E}_2^{p,q} = H^q(\mathbb{P}^d, \mathcal{H}^p(\mathcal{G}^\bullet)(m)), \quad \dot{d}_r^{p,q} : \dot{E}_r^{p,q} \longrightarrow \dot{E}_r^{p-r+1, q+r};$$

with common abutment $\mathbb{H}^{p+q}(\mathcal{G}^\bullet(m))$.

The term $\mathcal{G}^p(m)$ is a sum of line bundles of the form $\mathcal{O}_{\mathbb{P}^d}(m-\beta-1)$. Now $m-\beta-1 \geq m-d$ (this is clear from formula (12)), hence $H^{\geq 1}(\mathbb{P}^d, \mathcal{G}^p(m)) = 0$. Thus $E_1^{p,q} = 0$ for $q \geq 1$, this forces $E_2 = E_\infty$. Now $\dot{E}_2^{p,q}$ is zero outside the columns $p = 0, 1$, so $\dot{E}_3 = \dot{E}_\infty$.

Theorem 3.3. *Assume $m \geq 0$. Then*

$$\mathbb{H}^i = \begin{cases} H^0(\mathbb{P}^d, \mathcal{I}_X(m)) & \text{for } i = 0, \\ 0 & \text{for } i \neq 0, 1. \end{cases}$$

There is an extension

$$0 \longrightarrow H^1(\mathbb{P}^d, \mathcal{I}_X(m)) \longrightarrow \mathbb{H}^1 \longrightarrow \dot{E}_3^{1,0} \longrightarrow 0,$$

and $\dot{E}_3^{1,0}$ equals the image of the map

$$H^0(\mathbb{P}^d, f_*\mathcal{O}_Y(m)) \longrightarrow H^0(\mathbb{P}^d, \mathcal{D}(m)).$$

In particular, $\dot{E}_3^{1,0} = H^0(\mathbb{P}^d, \mathcal{D}(m))$ if $H^1(\mathbb{P}^d, \mathcal{O}_X(m)) = 0$.

We will use the following technical lemma in the proof.

Lemma 3.4. *For $m \geq 0$, the map $\dot{d}_2^{1,q} : H^q(\mathcal{D}(m)) \longrightarrow H^{q+2}(\mathcal{I}_X(m))$ is bijective for $q \geq 1$ and surjective for $q = 0$.*

PROOF. The map is a composite $H^q(\mathcal{D}(m)) \rightarrow H^{q+1}(\mathcal{O}_X(m)) \rightarrow H^{q+2}(\mathcal{I}_X(m))$ of two connecting homomorphisms in the long exact sequences associated to (6) and (7). (This follows directly from the definition of $\dot{d}_2$ (loc.cit.).) Now $H^{\geq 1}(\mathbb{P}^d, \mathcal{O}_{\mathbb{P}^d}(m)) = 0$, and we claim that $H^{\geq 1}(\mathbb{P}^d, f_*\mathcal{O}_Y(m)) = 0$. Indeed,

$$\begin{aligned} H^i(\mathbb{P}^d, f_*\mathcal{O}_Y(m)) &= H^i(Y, f^*\mathcal{O}_{\mathbb{P}^d}(m)) \quad (\text{by Leray spectral sequence}) \\ &= \bigoplus_{\sum i_r=i} \bigotimes_{r=1}^d H^{i_r}(\mathbb{P}^{e_r}, \mathcal{O}_{\mathbb{P}^{e_r}}(mr)) \quad (\text{by Künneth formula}) \end{aligned}$$

which vanishes for $i \geq 1$. It now follows that the first connecting map is a bijection for $q \geq 1$ and a surjection for $q = 0$; and the second map is always an isomorphism. The lemma is proved. $\square$

PROOF OF THEOREM 3.3.

Since $\dot{E}_2^{p,q} = 0$ for $p \neq 0, 1$, we have $\mathbb{H}^i = 0$ for $i < 0$. By the lemma, $\dot{E}_3^{p,q} = 0$ for $p = 0, q \geq 2$ and $p = 1, q \geq 1$; hence $\mathbb{H}^i = 0$ for $i \geq 2$. The rest is clear. $\square$

Corollary 3.5. *The space $H^0(\mathbb{P}^d, \mathcal{I}_X(m))$ of hypersurfaces of degree m vanishing on X is given by the middle cohomology of the complex*

$$H^0(\mathbb{P}^d, \mathcal{G}^{-1}(m)) \rightarrow H^0(\mathbb{P}^d, \mathcal{G}^0(m)) \rightarrow H^0(\mathbb{P}^d, \mathcal{G}^1(m)).$$

$\square$

At this point, the determination of this space is in principle a problem in linear algebra; once we have made the differential of the complex explicit. We will merely outline the description here and leave the details to the diligent reader.

Let $d_{G,m}^p : H^0(\mathbb{P}^d, \mathcal{G}^p(m)) \rightarrow H^0(\mathbb{P}^d, \mathcal{G}^{p+1}(m))$ denote the differential in question. Define

$$\begin{aligned} (16) \quad z(\alpha, r) &= r\alpha + r - e_r - 1, \quad M(\alpha) = \bigotimes_r \text{Sym}^{z(\alpha,r)}(\text{Sym}^{e_r} V) \quad \text{and} \\ Q(\alpha, p) &= M(\alpha) \otimes \text{Sym}^{m+p+\alpha-n-1}(\text{Sym}^d V^*) \otimes \wedge^{n+2-p}(\text{Sym}^d V^*). \end{aligned}$$

Then $H^0(\mathbb{P}^d, \mathcal{G}^p(m)) = \bigoplus_{n-p \geq \alpha \geq M-1} Q(\alpha, p)$, and it is only necessary to describe the differential $d_{G,m}^p : Q(\alpha, p) \rightarrow Q(\alpha', p+1)$ on individual summands. This map is zero unless $\alpha' = \alpha$ or $\alpha - 1$.

If $\alpha' = \alpha$, it suffices to describe the map

$$\begin{aligned} & \text{Sym}^{m+p+\alpha-n-1}(\text{Sym}^d V^*) \otimes \wedge^{n+2-p}(\text{Sym}^d V^*) \rightarrow \\ & \text{Sym}^{m+p+\alpha-n}(\text{Sym}^d V^*) \otimes \wedge^{n+1-p}(\text{Sym}^d V^*). \end{aligned}$$

This is given by contraction with the canonical trace element in $\text{Sym}^d V^* \otimes \text{Sym}^d V$. If $\alpha' = \alpha - 1$, it suffices to describe the map

$$M(\alpha) \otimes \wedge^{n+2-p}(\text{Sym}^d V^*) \rightarrow M(\alpha - 1) \otimes \wedge^{n+1-p}(\text{Sym}^d V^*).$$

This is given by contraction with the element in $\otimes_r \text{Sym}^r(\text{Sym}^{e_r} V^*) \otimes \text{Sym}^d V$, coming from the comultiplication map (9).

3.4. An Alternate Description of $\mathcal{I}_X$. We pick up the thread at the beginning of §3.2. There is a bundle map

$$(17) \quad \tilde{\varphi} : \pi^*(\Omega_{\mathbb{P}^d}^1(1)) \longrightarrow \mathcal{L}$$

to be defined as follows. (Let $\mathfrak{T}_{\mathbb{P}}$ denote the tangent bundle.) Specifying $\tilde{\varphi}$ is tantamount to specifying a global section in

$$H^0(T, \mathcal{L} \otimes \pi^* \mathfrak{T}_{\mathbb{P}^d}(-1)) = \bigotimes_{r=1}^d \text{Sym}^r(\text{Sym}^{e_r} V^*) \otimes \text{Sym}^d V,$$

which is tantamount to giving a map $\text{Sym}^d V^* \longrightarrow \bigotimes_{r=1}^d \text{Sym}^r(\text{Sym}^{e_r} V^*)$. Take this to be the comultiplication map (9).

Over the point $t = (G_1, \dots, G_d; F) \in T$, the transpose map $\mathcal{L}^{-1} \longrightarrow \pi^*(\mathfrak{T}_{\mathbb{P}^d}(-1))$ is given by

$$\tilde{\varphi}^{\text{trans}} : \prod_r G_r^r \longrightarrow \prod_r G_r^r + (F) \in \frac{\text{Sym}^d V}{(F)}.$$

The image is zero iff $F = \prod_r G_r^r$ upto constants, i.e., iff $t \in \Gamma$. Hence the scheme defined by $\text{Fitt}_0(\text{coker } \tilde{\varphi})$ is Γ . Now starting with the Koszul complex of $\tilde{\varphi}$, the arguments in §3.2, 3.3 go through verbatim. The outcome is a complex of bundles $\tilde{\mathcal{G}}^\bullet$ living on $\mathbb{P}^d$, with

$$(18) \quad \begin{aligned} \tilde{\mathcal{G}}^p &= \bigotimes_r \text{Sym}^{r(n+1-p)-e_r-1}(\text{Sym}^{e_r} V) \otimes \Omega_{\mathbb{P}^d}^{n+1-p}(n+1-p), \\ &\text{for } n+1-d \leq p \leq n+1-M. \end{aligned}$$

Then Theorem 3.3 and Corollary 3.5 are valid with $\tilde{\mathcal{G}}^\bullet$ in place of $\mathcal{G}^\bullet$.

Scholium 3.6. Consider the map

$$\gamma_m : \ker d_{G,m}^0 \longrightarrow H^0(\mathbb{P}^d, \mathcal{I}_X(m))$$

given by Theorem 3.3. The inclusion $\ker d_{G,m}^0 \subseteq H^0(\mathbb{P}^d, \mathcal{G}^0(m))$ is $SL(V)$ -equivariant, hence it must split (non-canonically). Hence γ_m extends to a map

$$\tilde{\gamma}_m : H^0(\mathbb{P}^d, \mathcal{G}^0(m)) \longrightarrow H^0(\mathbb{P}^d, \mathcal{I}_X(m)).$$

I believe that it would be of interest to have at least one explicit construction of $\tilde{\gamma}_m$.

4. SOME COMPUTATIONS

So far, we have described two approaches to the calculation $\mathcal{I}_X$, one in §3.1 and the other in Theorem 3.3. They can be combined to get a concrete invariant-theoretic description of this ideal, provided we can get a handle on $\mathcal{D}$.

We will carry out the calculation in detail for $\lambda = (3, 2)$, and then in addition give the result for (3, 3). In each case we will describe the minimal resolution of X_λ in terms of $SL_2(\mathbb{C})$ -representations. The theorem below will be used for (3, 2), but of course it is of more general application.

Theorem 4.1. *Let $\lambda = (\lambda_1, \lambda_2)$ be a partition of d , with $\lambda_1 \neq \lambda_2$. Then $\mathcal{D}_\lambda$ is supported on the rational normal curve $X_{(d)}$, and we have an isomorphism $\mathcal{D}_\lambda = f_{(d)*} \mathcal{O}_{\mathbb{P}^1}(-2)$.*

PROOF. The claim about $\text{supp}(\mathcal{D})$ will follow directly from Theorem 5.4 in the next section. To prove the rest, consider the commutative triangle

$$\begin{array}{ccc} \mathbb{P}^1 & \xrightarrow{\delta} & \mathbb{P}^1 \times \mathbb{P}^1 \\ & \searrow f_{(d)} & \downarrow f_\lambda \\ & & \mathbb{P}^d \end{array}$$

Here δ is the diagonal imbedding, let Δ be its image.

Step 1: The ideal of Δ equals $\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-1, -1)$, i.e., we have an extension

$$(19) \quad 0 \longrightarrow \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-1, -1) \longrightarrow \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1} \longrightarrow \mathcal{O}_\Delta \longrightarrow 0.$$

Consider the commutative ladder

$$\begin{array}{ccccccccc}
0 & \longrightarrow & \mathcal{I}_{X_{(d)}}/\mathcal{I}_{X_\lambda} & \longrightarrow & \mathcal{O}_{X_\lambda} & \longrightarrow & \mathcal{O}_{X_{(d)}} & \longrightarrow & 0 \\
& & \downarrow 1 & & \downarrow 2 & & \downarrow 3 & & \\
0 & \longrightarrow & f_{\lambda*}\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-1, -1) & \longrightarrow & f_{\lambda*}\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1} & \longrightarrow & f_{\lambda*}\mathcal{O}_\Delta & \longrightarrow & 0
\end{array}$$

whose bottom row is derived from (19). (The functor $f_{\lambda*}$ is exact since f_λ is a finite morphism.) Now the map 3 is an isomorphism, so by the five lemma, $\text{coker } 1 \cong \text{coker } 2$. This gives an extension

$$(20) \quad 0 \longrightarrow \mathcal{I}_{X_{(d)}}/\mathcal{I}_{X_\lambda} \xrightarrow{1} f_{\lambda*}\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-1, -1) \longrightarrow \mathcal{D}_\lambda \longrightarrow 0.$$

Step 2: There is an isomorphism $\delta^*\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-1, -1) \cong \mathcal{O}_{\mathbb{P}^1}(-2)$, hence by taking adjoints, a surjection $\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-1, -1) \rightarrow \delta_*\mathcal{O}_{\mathbb{P}^1}(-2)$. The kernel of this surjection equals $\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-2, -2)$, so we have an exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-2, -2) \longrightarrow \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-1, -1) \longrightarrow \delta_*\mathcal{O}_{\mathbb{P}^1}(-2) \longrightarrow 0.$$

After applying $f_{\lambda*}$, we have an extension

$$(21) \quad 0 \rightarrow f_{\lambda*}\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-2, -2) \rightarrow f_{\lambda*}\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-1, -1) \rightarrow f_{(d)*}\mathcal{O}_{\mathbb{P}^1}(-2) \rightarrow 0.$$

Step 3: From (20) and (21), it suffices to construct an isomorphism $\mathcal{I}_{X_{(d)}}/\mathcal{I}_{X_\lambda} \cong f_{\lambda*}\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-2, -2)$ to prove the theorem. It will be clear from the construction below that the isomorphism is compatible with the maps in (20) and (21).

Let $[x_0, x_1; y_0, y_1]$ (respectively $[a_0, \dots, a_d]$) be cöordinates on $\mathbb{P}^1 \times \mathbb{P}^1$ (respectively on $\mathbb{P}^d$). The morphism f_λ is defined via the identity

$$\sum_{j=0}^d a_j t^{d-j} = (x_0 t + x_1)^{\lambda_1} (y_0 t + y_1)^{\lambda_2}.$$

We pass to the open set $U_0 = \{a_0 \neq 0\}$ and write

$$\xi = x_1/x_0, \eta = y_1/y_0, \alpha_i = a_i/a_0 \text{ for } 1 \leq i \leq d.$$

The commutative triangle corresponds to a diagram of ring homomorphisms

$$\begin{array}{ccc}
 \frac{\mathbb{C}[\xi, \eta]}{(\xi - \eta)} & \xleftarrow{\text{epi}} & \mathbb{C}[\xi, \eta] \\
 & \nwarrow q_{(d)} & \uparrow q_\lambda \\
 & & \mathbb{C}[\alpha_1, \dots, \alpha_d]
 \end{array}$$

Let $I_\lambda = \ker q_\lambda$, $I_{(d)} = \ker q_{(d)}$.

Claim: The map $q_\lambda : I_{(d)} \longrightarrow \mathbb{C}[\xi, \eta]$ has image $(\xi - \eta)^2 \mathbb{C}[\xi, \eta]$. Hence it defines an isomorphism $I_{(d)}/I_\lambda \cong (\xi - \eta)^2 \mathbb{C}[\xi, \eta]$ of $\mathbb{C}[\alpha_1, \dots, \alpha_d]$ -modules.

Proof: The ideal $I_{(d)}$ has generators

$$J_r = (\alpha_1/d)^r - \alpha_r / \binom{d}{r} \quad \text{for } 2 \leq r \leq d.$$

Set $w = \xi - \eta$, then by straightforward computation,

$$\begin{aligned}
 q_\lambda(\alpha_1^r) &\equiv (d\eta)^r + r\lambda_1(d\eta)^{r-1}w, \\
 q_\lambda(\alpha_r) &\equiv \binom{d}{r}\eta^r + \lambda_1\eta^{r-1}\binom{d-1}{r-1}w \pmod{w^2};
 \end{aligned}$$

hence $q_\lambda(J_r) \equiv 0 \pmod{w^2}$. We will show that each term $w^2 \xi^i \eta^j$ lies in $q_\lambda(I_{(d)})$. Since $\lambda_2 \eta = q_\lambda(\alpha_1) - \lambda_1 \xi$, it suffices to show this for $j = 0$. Due to the equality

$$(\lambda_1^2 + \lambda_1 \lambda_2) \xi^2 = 2\lambda_1 q_\lambda(\alpha_1) \xi + (\lambda_2 - 1) q_\lambda(\alpha_1^2) - 2\lambda_2 q_\lambda(\alpha_2),$$

we may further assume $i = 0, 1$. Now a tedious calculation shows that a constant multiple of $q_\lambda(J_2)$ equals w^2 , and a certain linear combination of $q_\lambda(J_3)$ and $q_\lambda(\alpha_1 J_2)$ equals $w^2 \xi$. This proves the claim.

Evidently, the same calculation goes through over the open set $U_d = \{a_d \neq 0\}$, where we have an isomorphism

$$I_{(d)}/I_\lambda \cong (1/\xi - 1/\eta)^2 \mathbb{C}[1/\xi, 1/\eta].$$

The union $U_0 \cup U_d$ misses only two points of X_λ ; say P and Q , which are the images of points $[1, 0] \times [0, 1], [0, 1] \times [1, 0] \in \mathbb{P}^1 \times \mathbb{P}^1$. We have shown that over the open set $X_\lambda - \{P, Q\}$, we have an isomorphism $\mathcal{I}_{X_{(d)}}/\mathcal{I}_{X_\lambda} = f_{\lambda*} \mathcal{L}$ for some line bundle $\mathcal{L}$.

If g is an automorphism of X_λ coming from $SL(V)$, then $g^*(\mathcal{I}_{X_{(d)}}/\mathcal{I}_{X_\lambda}) \cong \mathcal{I}_{X_{(d)}}/\mathcal{I}_{X_\lambda}$. We can move P or Q to a point of $U_0 \cup U_d$ by such a g ,

so the isomorphism is valid throughout X_λ . Now take the ratio of the local generators

$$\frac{(x_1/x_0 - y_1/y_0)^2}{(x_0/x_1 - y_0/y_1)^2} = (x_0/x_1)^{-2}(y_0/y_1)^{-2},$$

which shows that $\mathcal{L} = \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-2, -2)$. The map 1 in (20) is the restriction of map 2, and locally, the latter is just q_λ at the level of rings. Hence the compatibility claimed earlier. This completes the proof of the theorem. $\square$

Corollary 4.2. *For λ as above, $H^0(\mathbb{P}^d, \mathcal{D}_\lambda(m)) = \text{Sym}^{dm-2} V^*$ for any $m \geq 0$. The Hilbert polynomial of X_λ is $\lambda_1 \lambda_2 m^2 + 2$.*

PROOF. We have $\mathcal{D}_\lambda \otimes \mathcal{O}_{\mathbb{P}^d}(m) = f_{(d)*} \mathcal{O}_{\mathbb{P}^1}(dm - 2)$ from the projection formula, hence the first claim. Now the second claim is a simple corollary of formula (7). $\square$

The following device will be useful for calculating with complexes of representations :

Let $\mathbb{A}$ be the free abelian group on symbols $\{s_n\}_{n \geq 0}$. We let $\chi(\text{Sym}^n V) = s_n$ for any n , and then define $\chi(R) \in \mathbb{A}$ for any finite dimensional $SL(V)$ -representation R by decomposing it into irreducibles. Thus $\chi(R)$ is merely the formal character of R .

If $G^\bullet$ is a bounded complex of $SL(V)$ -representations and $SL(V)$ -equivariant cochain maps, then define $\chi(G^\bullet) = \sum_p (-1)^p \chi(G^p)$. This can be seen as the ‘Euler characteristic’ of $G^\bullet$, taking values in $\mathbb{A}$. The following equality is immediate from Schur’s lemma :

$$(22) \quad \chi(G^\bullet) = \sum_p (-1)^p \chi(H^p(G^\bullet)).$$

4.1. The Surface X_{32} . As in §3.1, we have a morphism

$$f_{32} : \mathbb{P}V \times \mathbb{P}V \longrightarrow \mathbb{P}\text{Sym}^5 V, \quad (G_3, G_2) \longrightarrow G_3^3 \cdot G_2^2 (= F);$$

and in affine cöordinates, a ring map

$$q_{32} : \mathbb{C}[a_1, \dots, a_5] \longrightarrow \mathbb{C}[u, v],$$

$$a_1 \rightarrow 3u + 2v, \ a_2 \rightarrow 3u^2 + 6uv + v^2, \dots, \ a_5 \rightarrow u^3 v^2.$$

Homogenize the kernel of q_{32} with respect to a_0 , and take its minimal resolution:

$$\begin{aligned} 0 \longleftarrow \mathcal{I}_X \longleftarrow \mathcal{O}_{\mathbb{P}}(-4)^{28} \longleftarrow \mathcal{O}_{\mathbb{P}}(-5)^{68} \longleftarrow \mathcal{O}_{\mathbb{P}}(-6)^{64} \longleftarrow \mathcal{O}_{\mathbb{P}}(-7)^{28} \\ \longleftarrow \mathcal{O}_{\mathbb{P}}(-8)^5 \longleftarrow 0. \end{aligned}$$

All of this was carried out in Macaulay-2. We write this as

$$0 \longleftarrow \mathcal{I}_X \longleftarrow \mathcal{E}^0 \longleftarrow \mathcal{E}^{-1} \dots \longleftarrow \mathcal{E}^{-4} \longleftarrow 0.$$

Our task now is to identify the $SL(V)$ -representations entering into the syzygy modules $\mathcal{E}^p$. There is a spectral sequence

$$(23) \quad \begin{aligned} E_1^{p,q} &= H^q(\mathbb{P}^5, \mathcal{E}^p(m)), \quad d_r^{p,q} : E_r^{p,q} \longrightarrow E_r^{p+r, q-r+1} \\ E_\infty^{p,q} &\Rightarrow H^{p+q}(\mathbb{P}^5, \mathcal{I}_X(m)). \end{aligned}$$

Here $\mathcal{E}^p(m)$ denotes $\mathcal{E}^p \otimes \mathcal{O}_{\mathbb{P}^5}(m)$ etc. Note the equality

$$(24) \quad \begin{aligned} \chi(H^0(\mathcal{G}^\bullet(m))) &= \sum_p (-1)^p \chi(H^0(\mathbb{P}, \mathcal{G}^p(m))) \text{ by definition} \\ &= \chi(\mathbb{H}^0(\mathcal{G}^\bullet(m))) - \chi(\mathbb{H}^1(\mathcal{G}^\bullet(m))), \end{aligned}$$

which is merely (22) rewritten using Theorem 3.3.

Now let $m = 4$. From (23) we have $H^{\geq 1}(\mathcal{I}_X(4)) = 0$, so $H^1(\mathcal{O}_X(4)) = H^2(\mathcal{I}_X(4)) = 0$. (We are using nothing more than the standard computation of cohomology of line bundles on projective spaces, see [10, §3.5].) Now by Theorem 3.3,

$$\chi(\mathbb{H}^1(\mathcal{G}^\bullet(4))) = \chi(H^0(\mathbb{P}, \mathcal{D}(4))) = \chi(\text{Sym}^{18} V^*) = s_{18}.$$

The alternating sum $\chi(H^0(\mathcal{G}^\bullet(4)))$ can be calculated directly from (16) and repeated use of formulae (1) and (2). (This was programmed in Maple.) Substituting this into (24), we get

$$\chi(H^0(\mathcal{I}_X(4))) = \chi(\mathbb{H}^0(\mathcal{G}^\bullet(4))) = s_{12} + s_8 + s_4 + s_0.$$

Now choose $m = 5$. From (23), we deduce $H^{\geq 1}(\mathcal{I}_X(5)) = 0$ and an extension

$$0 \longrightarrow H^0(\mathcal{E}^{-1}(5)) \longrightarrow H^0(\mathcal{E}^0(5)) \longrightarrow H^0(\mathcal{I}_X(5)) \longrightarrow 0.$$

By the previous step,

$$H^0(\mathcal{E}^0(5)) = \text{Sym}^5 V^* \otimes (\text{Sym}^{12} V^* \oplus \text{Sym}^8 V^* \oplus \text{Sym}^4 V^* \oplus \mathbb{C}).$$

Now $\chi(H^0(\mathcal{G}^\bullet(5)))$ and $\chi(\mathbb{H}^1)$ are calculated exactly as before, and one can identify the unknown term

$$\chi(H^0(\mathcal{E}^{-1}(5))) = s_{13} + s_{11} + s_9 + 2s_7 + 2s_5 + s_3.$$

It is clear how to proceed to determine the entire minimal resolution. The complete result is as follows. (We have written e.g. $\{5, 3^2, 1^3\}$ for the module $\text{Sym}^5 V^* \oplus (\text{Sym}^3 V^*)^{\oplus 2} \oplus (V^*)^{\oplus 3}$.)

$$\begin{aligned} 0 \longleftarrow \mathcal{I}_X &\longleftarrow \{12, 8, 4, 0\} \otimes \mathcal{O}_{\mathbb{P}}(-4) \longleftarrow \{13, 11, 9, 7^2, 5^2, 3\} \otimes \mathcal{O}_{\mathbb{P}}(-5) \\ &\longleftarrow \{12, 10, 8^2, 6, 4^2, 2, 0^2\} \otimes \mathcal{O}_{\mathbb{P}}(-6) \longleftarrow \{9, 7, 5, 3\} \otimes \mathcal{O}_{\mathbb{P}}(-7) \\ &\longleftarrow \{4\} \otimes \mathcal{O}_{\mathbb{P}}(-8) \longleftarrow 0. \end{aligned}$$

It follows then, that $I_{X_{32}}$ is generated by a subrepresentation

$$(I_X)_4 = \text{Sym}^{12} \oplus \text{Sym}^8 \oplus \text{Sym}^4 \oplus \text{Sym}^0 \subseteq \text{Sym}^4(\text{Sym}^5 V^*).$$

We can identify the summands in the language of classical invariant theory. Define covariants

$$H = (F, F)^2, \quad i = (F, F)^4, \quad A = (i, i)^2$$

for the generic binary quintic F . (In classical literature the notation varies from source to source. I have consistently followed [7, Ch. VII], except that their f is our F . All the transvectants are calculated using the symbolic method, see Scholium 2.1.)

The inclusion $\text{Sym}^{12} \subseteq \text{Sym}^4(\text{Sym}^5 V^*)$ corresponds to a covariant of degree 4 and order 12, we tentatively call it J . (Said differently, let $J = \sum_i \varphi_i x^i y^{12-i}$. Then the vector space spanned by the forms φ_i is a subrepresentation of $(I_X)_4$, isomorphic to $\text{Sym}^{12} V^*$.) Now the space of covariants of type $(4, 12)$ is spanned by H^2 and $i.F^2$ (loc.cit.), hence $J = \alpha H^2 + \beta i.F^2$ for some constants α, β . Choose a form F lying in X_{32} (but otherwise sufficiently general) and evaluate the right hand side. By hypothesis J must vanish, from this we can deduce that $\alpha : \beta = 25 : -6$. Thus the inclusion $\text{Sym}^{12} \subseteq \text{Sym}^4(\text{Sym}^5 V^*)$ is entirely specified by the covariant $25 H^2 - 6 i.F^2$. The other summands in $(I_X)_4$ are calculated in the same way, and we get the following theorem:

Theorem 4.3. *A homogeneous binary quintic F can be factored as $L^3 M^2$ for some linear forms L, M ; if and only if, the covariants*

$$25 H^2 - 6 i.F^2, \quad 5 i.H + 6 F.(i, F)^2, \quad 2 i^2 + 15 (i, H)^2 \quad \text{and } A$$

vanish on F .

These are the ‘algebraic conditions’ on F , alluded to in the very beginning of the paper.

4.2. The surface X_{33} . We will merely state the result for the surface X_{33} , suppressing all computation. (This case is in fact easier than the previous one, since $\mathcal{D}_{33} = 0$.) The equivariant minimal resolution turns out to be

$$\begin{aligned} 0 \leftarrow \mathcal{I}_X &\leftarrow \{12, 8, 6\} \otimes \mathcal{O}_{\mathbb{P}}(-3) \leftarrow \{14, 12, 10^2, 8, 6^2, 4, 2^2\} \otimes \mathcal{O}_{\mathbb{P}}(-4) \\ &\leftarrow \{14, 12, 10^2, 8^2, 6^3, 4^2, 2\} \otimes \mathcal{O}_{\mathbb{P}}(-5) \leftarrow \{12, 10, 8^2, 6, 4^2, 2, 0^2\} \otimes \mathcal{O}_{\mathbb{P}}(-6) \\ &\leftarrow \{8, 6, 4\} \otimes \mathcal{O}_{\mathbb{P}}(-7) \leftarrow \{2\} \otimes \mathcal{O}_{\mathbb{P}}(-8) \leftarrow 0. \end{aligned}$$

Thus the ideal is generated in degree three by three covariants, and these can be identified as before. Again let $H = (F, F)^2$ and $i = (F, F)^4$ for the generic binary sextic F .

Theorem 4.4. *A homogeneous binary sextic F can be written as the cube of a quadratic form, if and only if, the covariants*

$$(F, H), (F, i) \text{ and } 8F \cdot (F, F)^6 - 75(F, i)^2$$

vanish on F .

It is almost superfluous to mention that Theorems 4.3 and 4.4 are not being seriously suggested as *practical criteria* for detecting multiple factors.

Scholium 4.5. Here we give the result for the case $\lambda = (3\ 2\ 2)$, which was left unfinished in §3.1. The sheaf $\mathcal{D}_{322}$ sits in an extension

$$0 \rightarrow f_{52*} \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-3, -1) \rightarrow \mathcal{D}_{322} \rightarrow f_{7*} \mathcal{O}_{\mathbb{P}^1}(-2) \rightarrow 0.$$

This is proved by an argument similar to (but more elaborate than) the one used in Theorem 4.1. From this, we deduce an exact sequence

$$0 \rightarrow \mathrm{Sym}^{5m-3} V^* \otimes \mathrm{Sym}^{2m-1} V^* \rightarrow H^0(\mathbb{P}^7, \mathcal{D}(m)) \rightarrow \mathrm{Sym}^{7m-2} V^* \rightarrow 0,$$

and now the calculation proceeds as before. The ideal is generated by the graded piece

$$\begin{aligned} (I_X)_6 &= \{30, 26, 24, 22^2, 20, 18^3, 16, 14^3, 12^2, 10^3, 8, 6^3, 2^2\} \\ &\subseteq \mathrm{Sym}^6(\mathrm{Sym}^7 V^*). \end{aligned}$$

Scholium 4.6. In the case $\dim X_\lambda = 1$ (i.e., of the rational normal curve) the minimal resolution is very pretty, so I cannot resist recording it here. Its ideal is generated by quadrics, more specifically

$$(I_X)_2 = \bigoplus_{r=1}^{[d/2]} \text{Sym}^{2d-4r} V^* \subseteq \text{Sym}^2(\text{Sym}^d V^*).$$

(The summands correspond to the transvectants $(F, F)^{2r}$ for $1 \leq r \leq [d/2]$.) The natural multiplication map

$$\text{Sym}^{d-1} V^* \otimes V^* \longrightarrow \text{Sym}^d V^* = H^0(\mathbb{P}^d, \mathcal{O}_{\mathbb{P}}(1)),$$

gives rise to a bundle map

$$\varphi : \mathcal{O}_{\mathbb{P}}(-1) \otimes \text{Sym}^{d-1} V^* \longrightarrow V \otimes \mathcal{O}_{\mathbb{P}}.$$

Then the inequality $\text{rank } \varphi \leq 1$ holds over a point $[w] \in \mathbb{P} \text{Sym}^d V$, if and only if, $w = v^d$ for some $v \in V$. Hence the Eagon–Northcott complex of φ resolves $\mathcal{O}_X$.

I enclose a list of representations entering into the minimal generators of I_X and the corresponding covariants when λ has two parts and $4 \leq d \leq 6$. For each d , define $H = (F, F)^2$, $i = (F, F)^4$. In each case I_m denotes the degree m part of I_X , the rest needs no explanation.

Case $d = 4$.

$$(31) : I_2 = \{0\} = \{i\}, I_3 = \{0\} = \{(F, H)^4\}.$$

$$(22) : I_3 = \{6\} = \{(F, H)\}.$$

Case $d = 5$.

$$(41) : I_2 = \{2\} = \{i\}.$$

$$(32) : I_4 = \{12, 8, 4, 0\} \\ = \{25 H^2 - 6 i \cdot F^2, 5 i \cdot H + 6 F \cdot (i, F)^2, 2 i^2 + 15 (i, H)^2, A\}.$$

Case $d = 6$.

$$(51) : I_2 = \{4, 0\} = \{i, (F, F)^6\}.$$

$$(42) : I_2 = \{0\} = \{(F, F)^6\}, I_3 = \{2\} = \{(F, i)^4\}, \\ I_4 = \{16, 12, 0\} = \{27 H^2 - 8 i \cdot F^2, 3 i \cdot H + 4 F \cdot (F, i)^2, (i, i)^4\}.$$

$$(33) : I_3 = \{12, 8, 6\} = \{(F, H), (F, i), 8 F \cdot (F, F)^6 - 75 (F, i)^2\}.$$

4.3. The Castelnuovo regularity of $\mathcal{I}_X$. Let $\mathcal{F}$ be a coherent $\mathcal{O}_{\mathbb{P}}$ -module. It is said to be m -regular for an integer m , if $H^i(\mathbb{P}, \mathcal{F}(m-i)) = 0$ for $i \geq 1$. If this is so, then $\mathcal{F}(m)$ is generated by global sections and $\mathcal{F}$ is m' -regular for any $m' \geq m$. This concept is due to David Mumford, see [14] for a masterly exposition.

Here we state a result bounding the regularity of $\mathcal{I}_{X_\lambda}$, when $\lambda = (\lambda_1, \lambda_2)$. Define

$$\mu = \begin{cases} d^2/4 - d + 3 & \text{for } \lambda_1 = \lambda_2, \\ \lambda_1 \lambda_2 (2\lambda_1 \lambda_2 - d + 2)^2 + 2\lambda_1 \lambda_2 - d + 5 & \text{otherwise.} \end{cases}$$

Theorem 4.7. *With λ as above, $\mathcal{I}_X$ is μ -regular. A fortiori, $\mathcal{I}_X$ is generated by forms of degree $\leq \mu$.*

PROOF. The first case follows directly from the central result of [13]. For the second case, note the following lemma.

Lemma 4.8 ([11],[14]). *Let $\mathcal{F}$ be a coherent $\mathcal{O}_{\mathbb{P}}$ -module and let H be a general hyperplane in $\mathbb{P}$. (It is sufficient that the generic point of H not be associated to $\mathcal{F}$.) Assume that $\mathcal{F}|_H$ is m -regular. Then $H^{\geq 2}(\mathbb{P}, \mathcal{F}(m-1)) = 0$ and $\mathcal{F}$ is $(h^1(\mathcal{F}(m-1)) + m)$ -regular.* $\square$

Now $\mathcal{I}_X|_H$ is the ideal of a nondegenerate curve of degree $2\lambda_1\lambda_2$ in $\mathbb{P}^{d-1}$; hence by [8, Theorem 1.1] it is m_0 -regular, where $m_0 = (2\lambda_1\lambda_2 - d + 3)$. From the surjection $H^0(\mathcal{O}_X(m_0 - 1)) \rightarrow H^1(\mathcal{I}_X(m_0 - 1)) \rightarrow 0$, we have $h^1(\mathcal{I}_X(m_0 - 1)) \leq h^0(\mathcal{O}_X(m_0 - 1))$. Since $H^{\geq 2}(\mathcal{I}_X(m_0 - 1)) = 0$, we deduce

$$h^0(\mathcal{O}_X(m_0 - 1)) = \chi(\mathcal{O}_X(m_0 - 1)) = \lambda_1 \lambda_2 (m_0 - 1)^2 + 2.$$

The theorem is proved. $\square$

It is not likely that the bound is optimal, see the last section for a conjecture. For any λ , if the Hilbert polynomial of $\mathcal{D}_\lambda$ is known, then such a bound can be formulated. See [11] for a general regularity bound along such lines².

²However, the definition of ‘ (b) -polynôme’ and the statement of ‘théorème principal’ are not very clear to me as stated there.

5. SINGULARITIES OF THE CR LOCUS

In case of a general partition λ , an explicit determination of $\mathcal{D}_\lambda$ on the lines of Theorem 4.1 does not appear to be easy. (The isomorphism $\mathcal{H}^1(\mathcal{G}^\bullet) = \mathcal{D}$ is of little help in this.) In this section, we solve the easier problem of describing the support of $\mathcal{D}_\lambda$, which coincides with the singular locus of X_λ .

Note that a point $F \in X_\lambda$ is nonsingular iff $\mathcal{O}_{X,F}$ is normal.

Proposition 5.1. *A point $F \in X$ is nonsingular if and only if the following conditions hold:*

1. *The preimage $f_\lambda^{-1}(F)$ is a singleton set, say $\{G\}$; and*
2. *the map $df_\lambda : T_{Y,G} \longrightarrow T_{\mathbb{P}^d,F}$ on tangent spaces is injective.*

PROOF. This is immediate from the local criterion of isomorphism in [9, Theorem 14.9]. $\square$

With the partition $\lambda = (1^{e_1} 2^{e_2} \dots d^{e_d})$, we now associate a set $\mathcal{S}_\lambda$ of partitions of d . The point of the definition lies in Theorem 5.4.

Definition 5.2. A partition $\mu = (1^{f_1} 2^{f_2} \dots d^{f_d})$ belongs to $\mathcal{S}_\lambda$ in one of the following (mutually exclusive) cases.

- a. There exist distinct integers r_1, r_2 such that μ is derived from λ by setting

$$\begin{aligned} f_{r_1} &= e_{r_1} - 1, & f_{r_2} &= e_{r_2} - 1, \\ f_{r_1+r_2} &= e_{r_1+r_2} + 1 \text{ and } f_r &= e_r \text{ elsewhere.} \end{aligned}$$

- b. There exist integers r_1, r_2, t such that $r_1 \neq r_2$, $e_{r_1} > 0$ and μ is derived from λ by setting

$$\begin{aligned} f_{r_1} &= e_{r_1} + 1, & f_{r_2} &= e_{r_2} - t \text{ and} \\ f_r &= e_r \text{ elsewhere.} \end{aligned}$$

(This forces $r_1 = t r_2$.)

- c. There exist integers r_1, r_2, r_3, t_1, t_2 such that r_1, r_2, r_3 are pairwise distinct, $r_3 = t_1 r_1 = t_2 r_2$, and μ is derived from λ by setting

$$\begin{aligned} f_{r_1} &= e_{r_1} - t_1, & f_{r_2} &= e_{r_2} - t_2, \\ f_{r_3} &= e_{r_3} + 2 \text{ and } f_r &= e_r \text{ elsewhere.} \end{aligned}$$

In each case λ is a refinement of μ , so $X_\mu \subseteq X_\lambda$. We will write $\mathcal{S}_\lambda = \mathcal{S}_\lambda^{(a)} \cup \mathcal{S}_\lambda^{(b)} \cup \mathcal{S}_\lambda^{(c)}$. Then $\mathcal{S}_\lambda$ is empty if and only if all parts in λ are equal.

Example 5.3. Let $\lambda = (1\ 2^3\ 3^2\ 4)$. Then $\mathcal{S}_\lambda^{(a)}$ has in all six elements, $(1\ 2^3\ 3\ 7), (1\ 2^2\ 3\ 4\ 5)$ etc. We can set $f_2 = 1, f_4 = 2$ and $f_r = e_r$ elsewhere, hence $(1\ 2^2\ 3^2\ 4^2) \in \mathcal{S}_\lambda^{(b)}$. We can set $f_6 = 2, f_3 = f_2 = 0$ and $f_r = e_r$ elsewhere, hence $(1\ 4\ 6^2) \in \mathcal{S}_\lambda^{(c)}$.

Theorem 5.4. *The singular locus of X_λ equals $\bigcup_{\mu} X_\mu$, the union quantified over all $\mu \in \mathcal{S}_\lambda$.*

The union may well be redundant, i.e., there may exist μ_1, μ_2 such that $X_{\mu_1} \subseteq X_{\mu_2}$. The theorem is a consequence of the following two propositions:

Proposition 5.5. *Let F be a singular point of X_λ .*

1. *If $f_\lambda^{-1}(F)$ is singleton, then $F \in X_\mu$ for some $\mu \in \mathcal{S}_\lambda^{(a)}$.*
2. *If $f_\lambda^{-1}(F)$ has more than one element and $F \notin X_\mu$ for any $\mu \in \mathcal{S}_\lambda^{(a)}$, then $F \in X_\mu$ for some $\mu \in \mathcal{S}_\lambda^{(b)} \cup \mathcal{S}_\lambda^{(c)}$.*

Proposition 5.6. *Let $F \in \bigcup_{\mu \in \mathcal{S}_\lambda} X_\mu$ be a point. Then either $f_\lambda^{-1}(F)$ is not singleton or $T_{Y,G} \longrightarrow T_{\mathbb{P}^d,F}$ is not injective for the unique G in $f_\lambda^{-1}(F)$. Hence in either case F is a singular point of X .*

The following lemma will be needed in the proofs.

Lemma 5.7. *Let $\mathbb{P} \operatorname{Sym}^m V \times \mathbb{P} \operatorname{Sym}^n V \xrightarrow{h} \mathbb{P} \operatorname{Sym}^{m+n} V$ denote the multiplication map $(A, B) \longrightarrow A.B (= C)$. The map*

$$dh : T_{\mathbb{P}^m \times \mathbb{P}^n, (A,B)} \longrightarrow T_{\mathbb{P}^{m+n}, C}$$

on tangent spaces is injective, if and only if A, B have no common factor.

PROOF. The map is given as

$$([\alpha_0, \dots, \alpha_m], [\beta_0, \dots, \beta_n]) \longrightarrow [\gamma_0, \dots, \gamma_{m+n}], \quad \gamma_k = \sum_{i+j=k} \alpha_i \beta_j.$$

By a linear change in x, y we may assume that $\alpha_0 = \beta_0 = 1$ at (A, B) . Then $\partial \gamma_k / \partial \alpha_i = \beta_{k-i}$, $\partial \gamma_k / \partial \beta_j = \alpha_{k-j}$, and the Jacobian matrix representing dh at (A, B) equals the Sylvester resultant of A and B . Hence it has rank $m + n$ exactly when A, B have no common factor. $\square$

Corollary 5.8. *If $G = (G_1, \dots, G_d) \in Y_\lambda$ is a point with $f_\lambda(G) = F$; then $df : T_{Y,G} \longrightarrow T_{\mathbb{P}^d,F}$ is injective if and only if no two of the G_r have a common factor.*

PROOF. The map Πv_r (see (5)) is injective on tangent spaces everywhere. Now write μ as a succession of maps with products of two factors at a time, and apply the lemma repeatedly. $\square$

In the sequel, we will use the same notation for two forms which differ by a nonzero multiplicative constant. Since there will be no occasion to add two forms, this should cause no confusion. If G is a form and L a linear form, then $\text{ord}_L(G) = m$ means that $L^m | G$, but $L^{m+1} \nmid G$.

PROOF OF PROPOSITION 5.5.

1. Say $f^{-1}(F) = \{G\}$, then by Prop. 5.1 and the previous corollary, two of the G_r have a common factor. Say a linear form L divides G_{r_1}, G_{r_2} . Define

$$\begin{aligned} Q_{r_1} &= G_{r_1}/L, & Q_{r_2} &= G_{r_2}/L, \\ Q_{r_1+r_2} &= L \cdot G_{r_1+r_2} \text{ and } Q_r &= G_r \text{ elsewhere.} \end{aligned}$$

Then Q defines a point of $f_\mu^{-1}(F)$ with $\mu \in \mathcal{S}_\lambda^{(a)}$. $\square$

2. Let $G = (G_r)$, $H = (H_r)$ be two distinct points in $f_\lambda^{-1}(F)$. No two of the G_r have a common factor (otherwise F would lie in X_μ for some $\mu \in \mathcal{S}_\lambda^{(a)}$) and similarly for the H_r . Let r_1 be the largest index such that $G_{r_1} \neq H_{r_1}$. (This of course means G_{r_1} is not a scalar multiple of H_{r_1} .) Then some linear factor L of G_{r_1} is not a factor of H_{r_1} . (Assume the contrary. Then some linear factor, say K , occurs with a higher power in G_{r_1} than in H_{r_1} . But then K must divide some $H_{r'}$ for $r' \neq r_1$. This is impossible since $H_{r_1}, H_{r'}$ have no common factors.) By the same argument, there exists a linear form M dividing H_{r_1} but not G_{r_1} . Say $\text{ord}_L(G_{r_1}) = \alpha$, $\text{ord}_M(H_{r_1}) = \beta$ and without loss of generality $\alpha \leq \beta$.

Case $\alpha = 1$. Now L divides $H_{r'}$ for exactly one value of r' , let r_2 be this value. If $\text{ord}_L(H_{r_2}) = t$, then $t r_2 = r_1$. Define

$$\begin{aligned} Q_{r_1} &= L \cdot H_{r_1}, & Q_{r_2} &= H_{r_2}/L^t \text{ and} \\ Q_r &= H_r \text{ elsewhere.} \end{aligned}$$

Then Q is a point of $f_\mu^{-1}(F)$ with $\mu \in \mathcal{S}_\lambda^{(b)}$.

Case $\alpha > 1$. Set $t_1 = \alpha$ and $r_3 = t_1 r_1$. Assume $\text{ord}_L(H_{r_2}) = t_2 > 0$ for some r_2 as above, then necessarily $r_2 t_2 = r_3$. Let

$$\begin{aligned} Q_{r_1} &= H_{r_1}/M^{t_1}, & Q_{r_2} &= H_{r_2}/L^{t_2}, \\ Q_{r_3} &= L.M.H_{r_3} \text{ and } Q_r &= H_r \text{ elsewhere.} \end{aligned}$$

Then Q is a point of $f_\mu^{-1}(F)$ with $\mu \in \mathcal{S}_\lambda^{(c)}$. $\square$

PROOF OF PROPOSITION 5.6. Assume that $F \in X_\mu$ for some $\mu \in \mathcal{S}_\lambda$ and that $f_\lambda^{-1}(F)$ is a singleton set $\{G\}$. We will deduce that two of the G_r must have a common factor; this will prove the proposition. In the sequel, H denotes a point of $f_\mu^{-1}(F)$.

Case $\mu \in \mathcal{S}_\lambda^{(a)}$. Choose a linear form L dividing $H_{r_1+r_2}$, and define

$$\begin{aligned} Q_{r_1} &= L.H_{r_1}, & Q_{r_2} &= L.H_{r_2}, \\ Q_{r_1+r_2} &= H_{r_1+r_2}/L \text{ and } Q_r &= H_r \text{ elsewhere.} \end{aligned}$$

Then $Q \in f_\lambda^{-1}(F)$, so $Q = G$. But then G_{r_1}, G_{r_2} have the common factor L .

Case $\mu \in \mathcal{S}_\lambda^{(b)}$. Note that $\deg H_{r_1} \geq 2$, and let L, L' be any two linear factors of H_{r_1} . We claim that $L = L'$. Suppose not, and define a point Q by

$$\begin{aligned} Q_{r_1} &= H_{r_1}/L, & Q_{r_2} &= H_{r_2}.L^t \text{ and} \\ Q_r &= H_r \text{ elsewhere;} \end{aligned}$$

and Q' by the same formulae with L' instead of L . Then $Q, Q' \in f_\lambda^{-1}(F)$ and $Q_{r_1} \neq Q'_{r_1}$, which is impossible. Hence necessarily $L = L'$, which implies that $L^2 | H_{r_1}$. But now $Q = G$, and Q_{r_1}, Q_{r_2} have the common factor L .

Case $\mu \in \mathcal{S}_\lambda^{(c)}$. Let L_1, L_2 be two linear factors of H_{r_3} and define

$$\begin{aligned} Q_{r_1} &= H_{r_1}.L_1^{t_1}, & Q_{r_2} &= H_{r_2}.L_2^{t_2}, \\ Q_{r_3} &= H_{r_3}/L_1 L_2 \text{ and } Q_r &= H_r \text{ elsewhere.} \end{aligned}$$

Define Q' by the same formulae with L_1, L_2 interchanged. Then $Q, Q' \in f_\lambda^{-1}(F)$, hence they must both equal G . Hence $L_1 = L_2$, and G_{r_1}, G_{r_2} have the common factor L_1 . $\square$

The proof of Theorem 5.4 is complete. On intuitive grounds it is plausible that the CR locus should develop a singularity wherever some parts in λ merge together. The element of surprise (for the author) lies in that only certain mergers produce singularities.

6. COMMENTS AND CONJECTURES

In his memoir [4], Cayley gives necessary and sufficient conditions for a form to lie in X_λ , for all partitions λ of $d \leq 5$. This amounts to describing I_X upto radicals. (Of course, his paper predates the recognition of the distinction between set-theoretic and ideal-theoretic generation.) E.g., for $\lambda = (3\,2)$, he proves that the summand $\text{Sym}^{12} \subseteq \text{Sym}^4(\text{Sym}^5 V^*)$ alone defines I_X set-theoretically³. However, *pace* Cayley's own assertion, I do not see how to make his method work for a general λ .

J. Weyman has proved [18, Theorem 3] that for $\lambda = (r\,1^{d-r})$ with $r \geq [d/2] + 1$, the ideal I_{X_λ} is generated in degrees ≤ 4 . (Also see [19] and [20] for results about the generators and Hilbert function of I_X .) I hazard the following conjecture (which I have numerically verified for $d \leq 8$).

Conjecture 6.1. Let $\lambda = (\lambda_1, \lambda_2)$ be a partition of d . If $\lambda_1 \neq \lambda_2$ (respectively $\lambda_1 = \lambda_2$), then the sheaf $\mathcal{I}_X(4)$ (resp. $\mathcal{I}_X(3)$) is Castelnuovo regular. In particular, I_X is generated in degrees ≤ 4 (resp. degrees ≤ 3).

It is proved in [15], that the variety $X_{(d-1,1)}$ is arithmetically Gorenstein. It is arithmetically Cohen–Macaulay in two more simple cases ($\lambda = (d), (2\,1^{d-2})$), but does not seem to be so for any other case. (I have numerically checked this for $d \leq 6$.) One would like to have a proof or a counterexample.

It would be valuable to have a structure theorem for $\mathcal{D}_\lambda$ similar to Theorem 4.1. There are a few more classes of partitions for which this can be achieved, and I hope to return to this problem in future.

REFERENCES

- [1] W. Adams and P. Loustaunau. *An introduction to Gröbner bases*. Graduate Studies in Mathematics, 3. American Mathematical Society, 1994.
- [2] E. Arbarello, M. Cornalba, P.A. Griffiths, and J. Harris. *Geometry of Algebraic Curves, Volume I*. Grundlehren der mathematischen Wissenschaften, No. 267. Springer–Verlag, New York, 1985.
- [3] R. Bott and L. Tu. *Differential Forms in Algebraic Topology*. Graduate Texts in Mathematics. Springer–Verlag, New York, 1982.

³Since Cayley's notation is different from ours, some effort is required to see that the first covariant in Theorem 4.3 agrees with his solution.

- [4] A. Cayley. A memoir on the conditions for the existence of given systems of equalities among the roots of an equation. *Philosophical Transactions of the Royal Society of London*, vol. cxlvii:pages 727–731, 1857. Reprinted in Volume II of his Collected Mathematical Papers by Cambridge Univ. Press, 1889.
- [5] D. Eisenbud. *Commutative Algebra, with a View Toward Algebraic Geometry*. Graduate Texts in Mathematics. Springer–Verlag, New York, 1995.
- [6] W. Fulton and J. Harris. *Representation Theory, A First Course*. Graduate Texts in Mathematics. Springer–Verlag, New York, 1991.
- [7] J. H. Grace and A. Young. *The Algebra of Invariants, 1903*. Reprinted by Chelsea, New York.
- [8] L. Gruson, R. Lazarsfeld, and C. Peskine. On a theorem of Castelnuovo, and the equations defining space curves. *Invent. Math.*, vol. 72, No. 3, pp. 491–506, 1983.
- [9] J. Harris. *Algebraic Geometry, A First Course*. Graduate Texts in Mathematics. Springer–Verlag, New York, 1992.
- [10] R. Hartshorne. *Algebraic Geometry*. Graduate Texts in Mathematics. Springer–Verlag, New York, 1977.
- [11] S. Kleiman. Les théorèmes de finitude pour le foncteur de Picard. In *SGA 6, Expose XIII*, Springer Lecture Notes in Mathematics No. 225, 1970.
- [12] J.P.S. Kung and G.-C. Rota. The invariant theory of binary forms. *Bulletin of the A.M.S.*, vol. 10, No. 1, pp. 27–85, 1984.
- [13] R. Lazarsfeld. A sharp Castelnuovo bound for smooth surfaces. *Duke Math J.*, vol. 55, No. 2, pp. 423–429, 1987.
- [14] D. Mumford. *Lectures on Curves on an Algebraic Surface*. Ann. of Math. Studies, No 59. Princeton University Press, 1966.
- [15] F.-O. Schreyer. Syzygies of canonical curves and special linear series. *Math. Ann.*, vol. 275, pp. 105–137, 1986.
- [16] T.A. Springer. *Invariant Theory*. Springer Lecture Notes in Mathematics No. 585, 1977.
- [17] B. Sturmfels. *Algorithms in Invariant Theory*. Texts and Monographs in Symbolic Computation. Springer–Verlag, Wien New York, 1993.
- [18] J. Weyman. The equations of strata for binary forms. *J. of Algebra*, vol. 122, No. 1, pp. 244–249, 1989.
- [19] J. Weyman. Gordan ideals in the theory of binary forms. *J. of Algebra*, vol. 161, No. 2, pp. 370–391, 1993.
- [20] J. Weyman. On Hilbert functions of multiplicity ideals. *J. of Algebra*, vol. 161, No. 2, pp. 358–369, 1993.

Department of Mathematics and Statistics,
 Queen’s University, Kingston, Ontario K7L 3N6. Canada.
 email: jaydeep@mast.queensu.ca